



PROCEEDINGS OF THE 11th WORLD RABBIT CONGRESS
Qingdao (China) - June 15-18, 2016
ISSN 2308-1910

Session
Management & Economy


**Szendrő K., Szendrő Zs., Gerencsér Zs.,
Radnai I., Horn P., Matics Zs.**

**ECONOMIC VALUE OF RABBIT LINES
SELECTED FOR DIFFERENT CRITERIA.**

**Full text of the communication
+
Slides of oral presentation**

How to cite this paper :

Szendrő K., Szendrő Zs., Gerencsér Zs., Radnai I., Horn P., Matics Zs., 2016 - Economic value of rabbit lines selected for different criteria. Proceedings 11th World Rabbit Congress - June 15-18, 2016 - Qingdao - China, 1013 - 1016 + Presentation.



ECONOMIC VALUE OF RABBIT LINES SELECTED FOR DIFFERENT CRITERIA

Szendró K.^{1*}, Szendró Zs.², Gerencsér Z.², Radnai I.², Horn P.², Matics Zs.²

¹ Faculty of Economic Science, Kaposvár University, 40 Guba S. St., H-7400, Kaposvár, Hungary

² Faculty of Agricultural and Environmental Sciences, Kaposvár University, 40 Guba S. St., H-7400, Kaposvár, Hungary

*Corresponding author: szendro.katalin@ke.hu

ABSTRACT

The aim of the experiment was to compare three genetic groups, slaughtered at similar weights, to examine their productive and carcass traits and economic value. Pannon Ka (PKa, maternal line) does were inseminated with semen of PKa, Pannon White (PWhite) or Pannon Large (PLarge, terminal line) bucks. The kits (PKa x PKa, PWhite x PKa, PLarge x PKa; n=60 in each genetic group) were weaned at 35 days of age and reared until 88, 83 and 79 days, respectively, when they reached similar body weights for slaughtering (2.8 kg). Results show that PLarge x PKa rabbits were able to exceed the average economic indicators compared to other groups. It can be concluded that the production performance of growing rabbits was affected by the adult weight, but the carcass traits were influenced by the computer tomography (CT)-based selection.

Key words: Growing rabbit, Lines, Production, Carcass traits, Economic value

INTRODUCTION

Most of the publications showed that rabbits originated from larger-sized parents (terminal lines) had better growth rate (Ramon *et al.*, 1996; Larzul and Rochambeau, 2004), but lower values of carcass traits (Dalle Zotte, 2002; Hernández *et al.*, 2006), since they were not as mature when slaughtered at the same weight as the progeny of maternal lines which had lower adult weights. Our hypothesis is that rabbits selected for L-value or TMV by computer tomography (CT) will be mature for slaughtering at younger age and achieve good slaughter results at a lower % of their adult weight than rabbits did not select for those traits.

The aim of the study was to compare economic values of three genetic groups slaughtered at similar weights to estimate whether the response in purebred within line selection – basically following a CTaimed procedure –, it is also observed under commercial condition.

MATERIALS AND METHODS

Animals and housing

The experiment was carried out at Kaposvár University. Three lines selected for different criteria, were examined in the experiment. Kaposvár University has a special breeding program, in which two lines (Pannon White and Pannon Ka) have been selected for a long time for carcass traits, based on the data of computer tomography (CT).. The characteristics of the lines of Pannon Breeding Program were summarized by Matics *et al.* (2014).

Pannon Ka (PKa) does (maternal line) were inseminated with semen of PKa, Pannon White (PWhite) or Pannon Large (PLarge) bucks. Randomly selected kits (PKa x PKa, PWhite x PKa, PLarge x PKa; n=60 in each genetic group) were weaned at 35 days of age and reared until 88, 83 and 79 days, respectively, when they reached similar body weights for slaughtering (2.8 kg). Rabbits were housed in a closed building in wire-mesh cages (3 rabbits/cage, 16 rabbits/m²). They were fed with commercial pelleted diets *ad libitum* and they could drink water freely from nipple drinkers. The temperature in the building was between 16 and 25 °C, and the daily lighting was 16 hours.

Measurements

Body weight and feed intake were measured from 5 weeks of age till they reached a similar body weight of 2.8 kg, therefrom weight gain was calculated. At the end of the experiment rabbits were slaughtered. The slaughtering and carcass dissection procedures followed the recommendations of World Rabbit Science Association (WRSA) described by Blasco and Ouhayoun (1996).

Financial indicators

Calculations are presented in Table 1. All financial figures were calculated in Euro. According to Maertens (2010) feeding cost may represent 60-70% of total production costs at the farm level. Since our experiment was carried out on growing rabbits, we used a value of 80% in calculating cost of production in case of medium price of feed (Maertens, personal communication). When calculating the cost of production on low and high feed price, – due to the change in the rate of feed cost – the percentage of feeding cost was 78.4 and 81.6%, respectively. Since cost of feed may vary significantly year by year, or even during a year and the market, the cost analysis was carried out based on the average cost of feed and 10% lower and 10% higher prices than the average price as well (low, medium /med/, high price).

The revenue from the whole rabbit carcass and from different carcass parts (revenue from rabbit products) was calculated. Based on these medium prices, 10% lower and 10% higher selling prices (low, med, high) were also calculated on the most valuable carcass parts (loin fillet and thigh meat), because the selling price of these items depends on different market prices. In these cases, a price change in whole carcass of +/- 8% was considered. Since the prices of other carcass parts (head, bones, fore part, etc.) are independent of the market, these were calculated on medium price

Table 1: Calculation of financial indicators (Cost and revenue based on farm and slaughterhouse level, €/rabbit)

Indicators	Cost of feed		
	Low (0.25 €/kg)	Med (0.275 €/kg)	High (0.3 €/kg)
1. Cost of weaned rabbit*		1.83 €/kg x body weight	
2. Cost of feeding**	Feed intake between weaning and slaughtering x cost of feed (+/-10%) (hay: 0.17 €/kg)		
3. Cost of mortality (dead rabbit)	1. + cost of feed (+/-10%) till death		
4. Cost of production	1. + 2. (78.4%) + 3.	1. + 2. (80%) + 3.	1. + 2. (81.6%) + 3.
5. Price at slaughter	1.53 €/kg* x body weight		
6. Farm profit	Revenue (price at slaughter) - Cost (Cost of production)		
7. Revenue from whole carcass*	Chilled carcass (g) x selling price (4.3 €/kg)		
8. Revenue from carcass parts*	[Loin fillet (12 €/kg); thigh meat (11 €/kg); liver (2.8 €/kg); kidneys (2.5 €/kg); fore part (2.6 €/kg); head, bone, heart and lungs (0.45 €/kg)] x weight of each carcass part		

*Data was gained from Olivia Ltd. ** Data were gained from Demeter /Cargill/

Statistical analysis

Statistical analysis was conducted using the SPSS 10.0 software package. The productive and carcass traits were evaluated by one-way ANOVA.

RESULTS AND DISCUSSION

Productive and carcass traits

Weight of PKa x PKa at weaning (889 g) was lower than that of the PLarge x PKa (923 g), and the largest in PWhite x PKa (947 g) groups ($P < 0.001$); presumably due to the differences in weight gain of kits between 3 and 5 weeks of age. Daily feed intake of PLarge x PKa rabbits (136 g/d) was significantly higher than that of PKa x PKa (127 g/d) and PWhite x PKa rabbits (129 g/d). These results were in accordance with those published in the literature (Ramon *et al.*, 1996; Feki *et al.*, 1996). However, the number of feeding days was less in the PLarge x PKa and more in PKa x PKa group, this is why the total feed consumption of PLarge x PKa was lower than that of PKa x PKa rabbits. Mortality was low, and no differences existed among the genetic groups.

In the present experiment rabbits were slaughtered at similar body weight. Despite finding no differences in body weight at slaughter, the weight of hind part, hind legs and meat on hind legs were higher in PWhite x PKa and PLarge x PKa groups, respectively, and the smallest in PKa x PKa rabbits ($P < 0.001$; table 2). The MLD was larger in PWhite x PKa and PKa x PKa rabbits than in group of PLarge x PKa ($P < 0.05$).

Table 3: Effect of genetic groups on carcass traits of rabbits (g) slaughtered at similar (2.8 kg) body weights

Traits	Genetic groups			SE
	PKa x PKa	PWhite x PKa	PLarge x PKa	
Age at the slaughter, d	88	83	79	
Body weight at slaughter	2785	2793	2795	9.7
Warm carcass	1708	1742	1726	6.9
Chilled carcass	1648	1678	1665	6.7
Fore part	418	418	425	2.2
Mid part	430	440	428	2.3
Hind part	512 ^a	537 ^b	530 ^b	2.5
Hind legs	476 ^a	501 ^b	495 ^b	2.4
HL	378 ^a	402 ^c	392 ^b	2.1
MLD	171 ^b	174 ^b	165 ^a	1.4

PKa = Pannon Ka (maternal line); PWhite = Pannon White; PLarge = Pannon Large (terminal line), HL = meat on hind legs; MLD = *m. longissimus dorsi*, ^{a,b,c}: Means in the same row with unlike superscripts differ at P<0.05 level.

Financial indicators

Cost of production and the profitability of different crossing combinations slaughtered at similar weights are shown in Table 4. The average difference in production costs (0.02 €/rabbit) was negligible between the PKa x PKa and PWhite x PKa groups, while a larger difference in cost of production (0.30 €/rabbit) was found between the former and PLarge x PKa in favor of PLarge x PKa rabbits, due to their shorter fattening period and, as a consequence, lower feed cost. Profit of PKa x PKa rabbits was 85 and 42% than that of the PWhite x PKa and PLarge x PKa group on a medium feed price, respectively. Results show that PLarge x PKa rabbits were able to exceed the average economic indicators on each feed price compared to the other groups.

A different ranking order occurred when the calculation was made at the slaughterhouse level. Revenues from rabbit carcass and rabbit products were 0.06 and 0.19, and 0.13 and 0.30 €/rabbit lower in PLarge x PKa and PKa x PKa rabbits, respectively, compared to PWhite x PKa group at a medium selling price. These differences were based on the carcass yields.

Results show a conflicting interest at farm and slaughterhouse level, since the farmer reached the highest benefit from PLarge x PKa, while the slaughterhouse in case of from PWhite x PKa rabbits, when they were sold and slaughtered at similar weight.

Table 4: Profitability of genetic groups (slaughtered at similar body weights, 2.8 kg) at the farm and slaughterhouse levels

Indicators	PKa x PKa			Genetic groups PWhite x PKa			PLarge x PKa		
	At farm								
	Price of feed								
	Low ¹	Med ¹	High ¹	Low ¹	Med ¹	High ¹	Low ¹	Med ¹	High ¹
Cost of feeding (€/r) ²	1.68	1.85	2.02	1.61	1.77	1.94	1.49	1.63	1.78
Cost of mortality (€/r) ²	0.09	0.09	0.09	0.06	0.06	0.06	0.00	0.00	0.00
Cost of production (€/r) ²	3.87	4.04	4.20	3.86	4.02	4.17	3.59	3.73	3.88
Price at slaughter (€/r) ²	4.27	4.27	4.27	4.28	4.28	4.28	4.29	4.29	4.29
Farm profit (€/r) ²	0.40	0.23	0.07	0.43	0.27	0.11	0.70	0.55	0.41
At slaughterhouse									
Selling price									
Low ¹	Med ¹	High ¹	Low ¹	Med ¹	High ¹	Low ¹	Med ¹	High ¹	High ¹
Revenue from rabbit carcass (€/r) ²	6.82	7.42	8.01	6.95	7.55	8.16	6.89	7.49	8.09
Revenue from rabbit products (€/r) ²	7.76	8.39	9.01	8.04	8.69	9.34	7.87	8.50	9.12

PKa = Pannon Ka (maternal line); PWhite = Pannon White; PLarge = Pannon Large (terminal line), ¹: Low, Med and High: low, medium and high price of pellets and selling price, ²: €/r= €/rabbit

CONCLUSIONS

It can be concluded that the production performance of growing rabbits was affected by the adult weight, but the carcass traits were influenced by the CT-based selection. The results of the present experiment showed new evidence, since the lines (PWhite and PLarge) which have been selected for carcass traits by CT had better dressing out percentage, ratio of hind part to reference carcass and profitability ratios than the maternal line (PKa) when they were compared at similar live weight range. The farmer reached the highest benefit from PLarge x PKa, while the slaughterhouse in case of PWhite x PKa rabbits when they were sold and slaughtered at similar weight. The Pannon Large as terminal line can improve the growth and also the carcass traits of progenies.

ACKNOWLEDGEMENTS

This paper was supported by the János Bolyai Research Scholarship of the Hungarian Academy of Sciences, and by the GOP-1.1.1-11-2012-0132 project

REFERENCES

- Dalle Zotte. A. 2002. Perception of rabbit meat quality and major factors influencing the rabbit carcass and meat quality. *Liv. Prod. Sci.*, 75, 11-32.
- Blasco A., Ouhayoun J. 1996. Harmonization of criteria and terminology in rabbit meat research. Revised proposal. *World Rabbit Sci.*, 4, 93-99.
- Feki S., Baselga M., Blas E., Cervera C., Gómez E.A. 1996. Comparison of growth and feed efficiency among rabbit lines selected for different objectives. *Liv. Prod. Sci.*, 45, 87-92.
- Hernández P., Ariño B., Grimal A., Blasco A. 2006. Comparison of carcass and meat characteristics of three rabbit lines selected for litter size or growth rate. *Meat Sci.*, 73, 645-650.
- Larzul C., de Rochambeau H. 2004. Comparison of ten rabbit lines of terminal bucks for growth, feed efficiency and carcass traits. *Anim. Res.*, 53, 535-545.
- Maertens L. 2010. Feeding systems for intensive production. In: De Blas C., Wiseman J. Eds, *Nutrition of the rabbit, 2nd Edition*. CAB International, Oxfordshire, 253-266.
- Matics Zs., Nagy I., Gerencsér Zs., Radnai I., Gyovai P., Donkó T., Dalle Zotte A., Curik I., Szendrő Zs. 2014. Pannon Breeding Program in rabbit at Kaposvár University. *World Rabbit Sci.*, 22, 287-300.
- Ramon J., Gómez E.A., Percho O., Rafel O., Baselga M. 1996. Feed efficiency and postweaning growth of several Spanish selected lines. In *Proc.: 6th World Rabbit Congress Toulouse, France*, 2, 351-353.
- SPSS for Windows 1999. Version 10.0. Copyright SPSS Inc.

=====



**K A P O S V Á R
U N I V E R S I T Y**

11TH WORLD RABBIT CONGRESS
JUNE 15-18, 2016, QINGDAO, CHINA



World Rabbit Congress
2016
Qingdao, China



ECONOMIC VALUE OF RABBIT LINES SELECTED FOR DIFFERENT CRITERIA

SZENDRŐ K.¹, SZENDRŐ Zs.², GERENCSÉR Zs.², RADNAI I.², HORN P.², MATICS Zs.²

¹Faculty of Economic Science, Kaposvár University, Hungary
²Faculty of Agricultural and Environmental Sciences, Kaposvár University, Hungary


szendro.katalin@ke.hu

INTRODUCTION




RABBITS ORIGINATED FROM LARGER-SIZED PARENTS (TERMINAL LINES, R) HAD BETTER GROWTH RATE, BUT LOWER VALUES OF CARCASS TRAITS (HIGHER FORE-, AND LOWER MID- AND HIND PARTS), SINCE THEY WERE NOT AS MATURE WHEN SLAUGHTERED AT THE SAME WEIGHT AS THE PROGENY OF MATERNAL LINES (P, V) WHICH HAD LOWER ADULT WEIGHTS.

Daily weight gain, g




Line	Daily weight gain, g
P	41.4
V	41
R	52

Body weight at 63 d, g



Line	Body weight at 63 d, g
P	2.09
V	2.12
R	2.6

Dressing out percentage, %



Line	Dressing out percentage, %
A	55.1
V	55.7
R	51.5

Source: Ramon et al., 1996; Larzul and Rochambeau, 2004; Dalle Zotte, 2002; Hernández et al., 2006; Pla, 1996



PRELUDE OF THE RESEARCH, AIM




**SELECTION FOR BETTER CARCASS TRAITS
BASED ON CT MEASUREMENT**

**K A P O S V Á R
U N I V E R S I T Y**



L-value **TMV**



**11-12 scans on the hind legs
(in each cm).**
**Pannon White were selected for
L-value since 1992.**
**Pannon White and Pannon Large
rabbits were selected for volume
of tight muscle (TMV) from 2004
and 2005, resp.**

**OUR HYPOTHESIS WAS THAT RABBITS SELECTED FOR L-VALUE OR TMV BY COMPUTER TOMOGRAPHY (CT)
WOULD BE MATURE FOR SLAUGHTERING AT YOUNGER AGE AND ACHIEVE GOOD SLAUGHTER RESULTS AT A
LOWER % OF THEIR ADULT WEIGHT THAN RABBITS DID NOT SELECT FOR THOSE TRAITS.**

PRELUDE OF THE RESEARCH, AIM

AIM

**K A P O S V Á R
U N I V E R S I T Y**

**The aim of the experiment was to compare
three genetic groups, slaughtered at similar
weights, to examine their productive and
carcass traits and economic value.**

PRELUDE OF THE RESEARCH, AIM

SOURCE OF DATA 



	4.0-4.5 kg	4.3-4.8 kg	4.8-5.4 kg
Adult body weight:	4.0-4.5 kg	4.3-4.8 kg	4.8-5.4 kg
Selected for:	<ul style="list-style-type: none"> •Litter size using BLUP method 	<ul style="list-style-type: none"> •daily weight gain (until 2010) •21d litter weight (from 2010) •L-value (until 2004) •thigh muscle volume (from 2004) 	<ul style="list-style-type: none"> •daily weight gain •thigh muscle volume (TMV)

 MATERIALS AND METHODS 

SOURCE OF DATA 



Pannon Ka (PKa)
Maternal line

Pannon White (PWhite)

Pannon Large (PLarge)

Kits (PKa x PKa, PWhite x PKa, PLarge x PKa; n=60 in each genetic group) were weaned at 35d and **reared until 88, 83 and 79 days, respectively, when they reached similar body weights (2.8 kg).**

 MATERIALS AND METHODS 




TABLE 1: FINANCIAL INDICATORS AT FARM LEVEL (€/rabbit)



Indicators	Cost of feed		
	Low 0.25 €/kg	Med 0.275 €/kg	High 0.3 €/kg
1. Cost of weaned rabbit*	1.83 €/kg x body weight		
2. Cost of feeding**	Feed intake between weaning and slaughtering x cost of feed (+/-10%)		
3. Cost of mortality (dead rabbit)	1. + cost of feed (+/-10%) till death		
4. Cost of production	1. + 2. (78.4 80 +81.6%) + 3.		
5. Price at slaughter	1.53 €/kg* x body weight		
6. Farm profit	Revenue (price at slaughter) - Cost of production		

*Data was gained from Olivia Ltd. ** Data were gained from Demeter /Cargill/



MATERIALS AND METHODS




**TABLE 2: FINANCIAL INDICATORS
AT **SLAUGHTERHOUSE** LEVEL (€/rabbit)**



Indicators	Selling price		
	Low	Med	High
1. Price at slaughter	1.53 €/kg* x body weight		
2. Revenue from whole carcass*	Chilled carcass (g) x selling price (4.3 €/kg)		
3. Revenue from carcass parts*	[Loin fillet (12 €/kg); thigh meat (11 €/kg); liver (2.8 €/kg); kidneys (2.5 €/kg); fore part (2.6 €/kg); head, bone, heart and lungs (0.45 €/kg)] x weight of each carcass part		

*Data was gained from Olivia Ltd.



MATERIALS AND METHODS



**TABLE 3: EFFECT OF GENETIC GROUPS ON CARCASS TRAITS OF
RABBITS (G) SLAUGHTERED AT SIMILAR (2.8 KG) BODY WEIGHTS**

Traits	Genetic groups			SE
	PKa x PKa	PWhite x PKa	PLarge x PKa	
Age at the slaughter, d	88	83	79	
Body weight at slaughter	2785	2793	2795	9.7
Warm carcass	1708	1742	1726	6.9
Chilled carcass	1648	1678	1665	6.7
Fore part	418	418	425	2.2
Mid part	430	440	428	2.3
Hind part	512 ^a	537 ^b	530 ^b	2.5
Hind legs	476 ^a	501 ^b	495 ^b	2.4
HL meat	378 ^a	402 ^c	392 ^b	2.1
MLD meat	171 ^b	174 ^b	165 ^a	1.4


PKa = Pannon Ka (maternal line); PWhite = Pannon White; PLarge = Pannon Large (terminal line), HL = meat on hind legs; MLD = *m. longissimus dorsi*,

^{a,b,c}: Means in the same row with unlike superscripts differ at P<0.05 level.



RESULTS





CONCLUSIONS

▪ The **farmer** reached the highest benefit from PLarge x Pka rabbits.

▪ The **slaughterhouse** reached the highest benefit in case of PWhite x PKa rabbits.

Production performance of growing rabbits was affected by the adult weight, but the **carcass traits** were influenced by the CT-based selection. The lines (PWhite and PLarge) which have been selected for carcass traits by CT had better dressing out percentage, ratio of hind part to reference carcass and profitability than the maternal line (PKa) when they were compared at similar live weight range.

CONCLUSIONS

Thank you for your
kind attention!



This paper was supported by the János Bolyai Research Scholarship of the Hungarian Academy of Sciences,
and by the GOP-1.1.1-11-2012-0132 project