



## **PROCEEDINGS OF THE 12<sup>th</sup> WORLD RABBIT CONGRESS**

Nantes (France) - November 3-5, 2021

ISSN 2308-1910

Session

**BIOLOGY and PHYSIOLOGY**

***Mattioli S., Cartoni Mancinelli A., Cotozzolo E., Mancini S.,  
Castellini C., Dal Bosco A.***

**COMPARISON OF AN ESTIMATED INDEX OF FATTY ACID METABOLISM AND  
LIVER  $\Delta$ 6-DESATURASE ACTIVITY IN RABBIT**

**Full text of the communication**

+

**Poster**

### *How to cite this paper*

Mattioli S., Cartoni Mancinelli A., Cotozzolo E., Mancini S., Castellini C., Dal Bosco A., 2021. Comparison of an estimated index of fatty acid metabolism and liver  $\Delta$ 6-desaturase activity in rabbit. Proceedings 12th World Rabbit Congress - November 3-5 2021 - Nantes, France, Communication BP-29, 4pp. + presentation

## COMPARISON OF AN ESTIMATED INDEX OF FATTY ACID METABOLISM AND LIVER $\Delta 6$ -DESATURASE ACTIVITY IN RABBIT

Mattioli S.\*<sup>1</sup>, Cartoni Mancinelli A.<sup>1</sup>, Cotozzolo E.<sup>1</sup>,  
Mancini S.<sup>2</sup>, Castellini C.<sup>1</sup>, Dal Bosco A.<sup>1</sup>

<sup>1</sup>Department of Agricultural, Food and Environmental Sciences, University of Perugia, 06121 Perugia, Italy

<sup>2</sup>Department of Veterinary Sciences, University of Pisa, 56124 Pisa 56124, Italy.

\*Corresponding author: simona.mattioli@unipg.it

### ABSTRACT

The aim of this study was to verify the accuracy of  $\Delta 6$ -desaturase ( $\Delta 6D$ ) index estimated from fatty acid profile of liver and meat of rabbit in comparison with its real metabolic activity directly evaluated in the liver microsomes. Two experiments were carried out using different diets (Control vs Linolenic diets) and genotypes (selected vs local breeds) for the comparison of  $\Delta 6D$  index and enzyme activity assay. Concerning the feed effect, a lower estimated index was found in Linolenic samples, both in liver and muscle, and this trend was only confirmed by the desaturase activity towards the n-3 fatty acids. The total and n-6 fatty acid-enzyme activity did not show significant differences between groups, as well as any correspondences with the indexes. The “selected” rabbits showed lower indexes respect to the local ones. The activity of the enzymatic complex has instead highlighted an opposite trend for the n-3 and the n-6 series and total one. In particular, the enzyme activity toward n-3 followed the same tendency of the index, whereas the other two determinations showed an opposite trend. The results of this comparison demonstrate that the calculation of the proposed index could be used as surrogates of the measure of liver  $\Delta 6$ -desaturase activity in lipid metabolism of rabbit, but only if specific n-3 pathway is considered.

**Key words:** Rabbit, meat, lipid metabolism, desaturase, fatty acids

### INTRODUCTION

In recent years, the fatty acid composition of meat has been intensively studied considering its strong implications in human health. Meat of monogastric animals could represent a source of long-chain n-3 polyunsaturated fatty acids (n-3LCP) conversely to the fish, whose consumption is generally low in European countries (Welch et al., 2002). Fish and livestock animals have developed different abilities to convert the precursor C18:3n-3 ( $\alpha$ -Linolenic acid, LNA) into n-3LCP, greatly depending on the efficiency of elongase and desaturase enzymes (mainly represented by the  $\Delta 6$ -desaturase complex -  $\Delta 6D$ ). Therefore, the assessment of activity, expression and substrate preference of this enzymatic complex is becoming of a certain importance in animal science in order to deepen the knowledge on lipid metabolism and to ameliorate the nutritional characteristics of the livestock products. Being the determination of the activity of these enzymes laborious and expensive, some Authors (Vessby et al., 2002) proposed certain indexes as surrogates of the measure of the “true” elongase/desaturase activity. The activity of  $\Delta 6D$  is downregulated by high dietary LCP fatty acids level, and hence, the degree of conversion of essential fatty acids to their longer and more unsaturated metabolites is lower (Nakamura et al., 2004; Rodríguez et al., 2019). Moreover, genetic selection can modify the expression of genes coding for enzymes involved in LCP synthesis as well as the relative enzymatic activity (Castellini et al., 2016).

To the best of our knowledge, no comparison between estimation and analytical determination have been carried out in rabbit; thus, the aim of the present study was to define the real applicability of an estimated index of fatty acid metabolism, through its comparison with the  $\Delta 6D$  enzyme activity assay.

## MATERIALS AND METHODS

### Animals and experimental design

#### Experiment 1

Forty male New Zealand White rabbits at the weaning age (30 days) were housed in the experimental farm of the University of Perugia (temperature 18-23 °C, humidity 50 ± 5%) and divided in two homogeneous groups. Each treatment consisted of 20 rabbits and each group receiving a different diet (Table 1) as follows:

- Control group was fed *ad libitum* with the control diet containing soya bean oil as the main source of fat;
- Linolenic group was fed the control diet, substituting 10% of the soy bean oil with extruded flaxseed;

**Table 1.** Formulation (%) of experimental diets.

Ingredients	Control	Linolenic
Barley meal	9.0	5.0
Fine wheat bran	30.0	26.4
Dehydrated alfalfa meal	40.0	50.00
Soybean meal 44% crude protein	4.0	3.50
Soybean oil	0.5	-
Sunflower meal 30% crude protein	9.0	5.00
Extruded linseed	3.0	10.0
Estimated digestible Energy**(MJ)	2.34	2.38

#### Experiment 2

Forty males (30 days) rabbits of two different genotypes were used and reared as in Experiment 1. In particular, both twenty commercial rabbits (New Zealand White rabbits) and rabbits of local strain (Grey-colored Italian rabbits, conserved for more than 30 years in small-sized farms in Central Italy) were used and fed with the above described control diet.

### Sampling and analysis

At 85 d, 10 rabbits per group of the two experiments, with weights close to the average of the group ( $\pm 10\%$ ), were selected and killed by cutting their carotid arteries and jugular veins after electrical stunning. Handling and dissection of the refrigerated carcasses (24 h at 4 °C) were performed as proposed by Blasco and Ouhayoun (1996). Within a few minutes, samples of liver and *Longissimus lumborum* (LL) muscle were cut into appropriately sized pieces (30 mg). Livers samples were partially rinsed with RNase-free water and frozen at -80 °C for subsequent evaluation of enzyme activity (Maranesi et al., 2015). Lipid extraction of liver and *Longissimus lumborum* muscle was performed according to Folch et al. (1957) and esterification was performed according to Christie (1982). The trans-metilation procedure was carried out using enicosenoic acid methyl esters (Sigma Chemical Co.) as the internal standard. The fatty acid composition was determined using a Varian gas-chromatograph (CP-3800) equipped with a flame ionisation detector and a capillary column of 100 m length  $\times$  0.25mm  $\times$  0.2  $\mu$ m film (Supelco, Bellefonte, PA, USA). Helium was used as the carrier gas with a flow of 2 mL/min.

The  $\Delta 6$ -desaturase ( $\Delta 6D$ ) activity was determined for the different PUFA series (n-6 and n-3) by measuring respectively, the amounts of 18:3n-6 produced from 18:2n-6 (Linoleic acid, LA) and 18:4n-3 produced from LNA. The reaction medium, in a total volume of 0.5 mL, contained the following: 4 mM ATP, 0.5 mM CoASH, 1.25 mM NADH, 2 mM MgCl<sub>2</sub>, 12 mM NaF, 1.5 mM glutathione, 0.5 M potassium phosphate, pH 7.2, 40 nmol 1-<sup>14</sup>C-labeled fatty acids (AS  $\cong$  3 nCi/nmol), and 5mg of microsomal proteins. Considering the amount of free LA and LNA in microsomes, in LA incubations totals of about 70 nmol and 64 nmol LA were present, whereas endogenous LNA was almost negligible

(about 4 nmol in F and 8 nmol in S). In LNA incubations, the activity was determined in the presence of about 30 nmol (F) and 24 nmol (S) LA.

To estimate the activity of  $\Delta 6$ -desaturase the following index was calculated from the fatty acid profile of liver and muscle (Dal Bosco et al., 2012):

$$\frac{[(C20:2n-6 + C20:4n-6 + C20:5n-3 + C22:5n-3 + C22:6n-3)]}{(LA + LNA + C20:2n-6 + C20:4n-6 + C20:5n-3 + C22:5n-3 + C22:6n-3)} \times 100.$$

### Statistical analysis

A linear model (STATA, 2005, procedure GLM) was used to evaluate the effect of feeding and genotype on estimated and real  $\Delta 6$ -desaturase activity. The statistical significance of differences was assessed using t-tests.

## RESULTS AND DISCUSSION

The estimated indexes and enzyme activity of  $\Delta 6$ -desaturase in rabbits fed different diets or belonging to two different genotypes are reported in Table 2.

**Table 2** Estimated and enzyme activity of  $\Delta 6$ -desaturase (pmol in 30 min/mg protein) in rabbits fed different diets or belonging to different genotypes

	Diets		Pooled SE <sup>1</sup>
	Control	Linolenic	
Estimated $\Delta 6$ -desaturase index (liver)	36.2 <sup>b</sup>	28.9 <sup>a</sup>	2.4
Estimated $\Delta 6$ -desaturase index (LL muscle)	28.6 <sup>b</sup>	25.3 <sup>a</sup>	2.4
$\Delta 6$ -desaturase for C18:3n-3	20.5 <sup>b</sup>	17.2 <sup>a</sup>	3.8
$\Delta 6$ -desaturase for C18:2n-6	177.5	181.2	10.8
Total $\Delta 6$ -desaturase activity	198.0	198.4	14.7
	Genotypes		Pooled SE <sup>1</sup>
	Commercial	Local	
Estimated $\Delta 6$ -desaturase index (liver)	31.8 <sup>a</sup>	33.1 <sup>b</sup>	1.5
Estimated $\Delta 6$ -desaturase index (LL muscle)	27.8 <sup>a</sup>	30.6 <sup>b</sup>	1.5
$\Delta 6$ -desaturase for C18:3n-3	19.1 <sup>a</sup>	26.9 <sup>b</sup>	2.4
$\Delta 6$ -desaturase for C18:2n-6	175.2 <sup>b</sup>	41.5 <sup>a</sup>	15.0
Total $\Delta 6$ -desaturase activity	194.3 <sup>b</sup>	68.4 <sup>a</sup>	17.5

Means in the same row and tissue with different letters differ significantly ( $P \leq 0.05$ ).

<sup>1</sup> SE: Standard Error

Concerning the feeding effect, lower estimated indexes were calculated in Linolenic group respect to Control (both in liver and muscle), and this trend was confirmed by the enzyme activity towards n-3. On the contrary, the total and n-6 oriented  $\Delta 6$ D activity did not show significant differences between groups. It is widely known that n-6 and n-3 fatty acid series compete for the same enzymes, but the affinity of the  $\Delta 6$ D is retained greater for n-3 than for n-6 fatty acids. Thus, the conversion of LA to n-6LCP metabolites decreases when the intake of n-3 fatty acids increases (Lands, 1992). Nevertheless, the above-mentioned preference of  $\Delta 6$ D for n-3 is neutralised by the fact that standard diet had higher content of n-6 fatty acids, especially LA, respect to Linolenic. The calculation of the indexes in different genotypes confirmed our previous findings (Dal Bosco et al., 2014) and in particular, that the fatty acid metabolism of animals selected for productive performance differed from those of local strains. Specifically, the NZW rabbits always showed lower indexes respect to the local ones.

The results of the present study confirm our previous study in poultry, where the estimated enzyme  $\Delta 6$ D activity was always higher in local breeds than in much selected commercial hybrids (Dal Bosco et al., 2012). Hence, the higher estimated  $\Delta 6$ D observed in local rabbits, can suggest a genetic effect on the

mechanisms of desaturase and elongase responsible for the synthesis of LCP fatty acids. Moreover, the real activity of the  $\Delta 6D$  enzymatic complex highlighted an opposite trend for the n-3 than the n-6 series and total activity. In particular, the enzyme activity for n-3 fatty acids followed the same trend of the estimated indexes, whereas in the other two determinations the trend was opposite. These results agree with Castellini et al. (2016), which observed a lower total  $\Delta 6D$  activity in slow growing rabbits respect to fast growing ones (45.2 vs 104.4 pmol/30 min/mg protein). In the latter research about 90 % of the activity was directed toward LA, whereas only 10 % toward LNA. In contrast, the  $\Delta 6D$  activity of slow-growing rabbits was 38.5% toward LNA and 61.5 % toward LA with higher affinity for LNA. These results showed that genetic selection for productive traits in rabbit modifies the expression and the activity of  $\Delta 6D$ . Thereafter, the  $\Delta 6$ -desaturase activity seems affected by precursor availability, but the major modifications in the LA/LNA ratio do not change substantially this tendency and that preference for LCP series in rabbit is mainly due to the intensity of genetic selection.

## CONCLUSIONS

The results of this comparison represent a first step of research towards a hypothetical use of the proposed index in the study of the rabbit lipid metabolism as surrogate of the true desaturase/elongase activity. In this study, the  $\Delta 6D$  index only mimic the activity toward the n-3 series would seem usable; indeed; indeed, the trend of estimated indexes and the total and n-6 direct enzyme activities were not superimposable. Obviously, further and thorough studies are needed both to expand the available database and to deepen the knowledge on the lipid metabolism in rabbit, and finally, to validate the potential applicability of the indexes.

## ACKNOWLEDGEMENTS

The research was funded by ‘Ricerca di base 2018’ University of Perugia

## REFERENCES

- Blasco A., Ouhayoun J. 1996. Harmonization of criteria and terminology in rabbit meat research. Revised proposal. *World Rabbit Sci.*, 4: 93-98.
- Castellini C, Dal Bosco A, Mattioli S, Davidescu M, Corazzi L, Macchioni L, Rimoldi S and Terova G. 2016. Activity, expression, and substrate preference of the  $\delta 6$ -desaturase in slow- or fast-growing rabbit genotypes. *J. Agric. Food Chem.*, 64: 792-800.
- Christie W.W. 1982. A simple procedure for rapid transmethylation of glycerolipids and cholesteryl esters. *J. Lipid Res.*, 23: 1072-1075.
- Dal Bosco A., Mugnai C., Ruggeri S., Mattioli S., Castellini C. 2012. Fatty acid composition of meat and estimated indices of lipid metabolism in different poultry genotypes reared under organic system. *Poultry Sci.*, 91: 2039-2045.
- Dal Bosco A., Mugnai C., Roscini V., Paci G., Castellini C. 2014. Effect of genotype on estimated indexes of fatty acid metabolism in rabbits. *World Rabbit Sci.*, 22: 21-28.
- Folch J., Lees M., Sloane-Stanley H. 1957. A simple method for the isolation and purification of total lipids from animal tissue. *J. Biol. Chem.*, 226: 497-509.
- Lands W. E. 1992. Biochemistry and physiology of n-3 fatty acids. *The FASEB journal*, 6(8), 2530-2536.
- Maranesi M., Zerani M., Leonardi L., Pistilli A., Arruda-Alencar J., Stabile A.M., Rende M., Castellini C., Petrucci L., Parillo F., Moura A., Boiti C. 2015. Gene expression and localization of NGF and its cognate receptors NTRK1 and NGFR in the sex organs of male rabbits. *Repr. Dom. Anim.* 50: 918-925.
- Nakamura MT and Nara TY 2004. Structure, function, and dietary regulation of delta6, delta5, and delta9 desaturases. *Annual Rev.Nutr.*, 24:345-376.
- Rodríguez, M., G Rebollar, P., Mattioli, S., Castellini, C. 2019. n-3 PUFA Sources (Precursor/Products): A Review of Current Knowledge on Rabbit. *Animals*, 9(10), 806.
- StataCorp. 2005. Stata Statistical Software: Release 9.0, College Station, Texas, USA.
- Vessby B., Gustafsson I.B., Tengblad S., Boberg M., Andersson A. 2002. Desaturation and elongation of fatty acids and insulin action. *Ann. NY Acad. Sci.*, 67: 183-195.
- Welch A.A, Lund E., Amiano P., Dorransoro M., Brustad M., Kumle M., Rodriguez M., Lasheras C., Janzon L., Jansson J., Luben R., Spencer E.A., Overvad K., Tjønneland A., Clavel-Chapelon F., Linseisen J., Klipstein-Grobusch K., Benetou V., Zavitsanos X., Tumino R., Galasso R., Bueno-de-Mesquita H.B., Ocké M.C., Charrondière U.R., Slimani N. 2002. Variability of fish consumption within the 10 European countries participating in the European Investigation into Cancer and Nutrition (EPIC) study. *Public Health Nutr.*, 5: 1273-1285.

# Comparison of an estimated index of fatty acid metabolism and liver $\Delta 6$ -desaturase activity in the rabbit



Mattioli S.<sup>1\*</sup>, Cartoni Mancinelli A.<sup>1</sup>, Cotozzolo E.<sup>1</sup>,  
Mancini S.<sup>2\*</sup>, Castellini C.<sup>1</sup>, Dal Bosco A.<sup>1\*</sup>

Email: simona.mattioli@unipg.it

\* attending authors

<sup>1</sup>Department of Agricultural, Food and Environmental Sciences, University of Perugia, Italy

<sup>2</sup>Department of Veterinary Sciences, University of Pisa, Italy.

**Context:** It is possible to hypothesize the use of an estimated index of lipid metabolism **as surrogate** of the determination of **true** desaturase/elongase activity in rabbit?

## Methods:

### Exp.1: feed effect

Control group

Linolenic group + 10% of flax

### Exp.2: strain effect

New Zealand White

Grey-colored Italian



From fatty acid composition of liver and muscle index of  $\Delta 6$ -desaturase activity ( $\Delta 6D$ ) according to Dal Bosco et al. (2012) was estimated; The real  $\Delta 6D$  activity was determined in liver microsomes.

## Results:

Lower estimated indexes were observed in Linolenic group as well as the enzyme activity towards n-3 (**RED**).

Regarding the genotype effect, the same overlap, albeit increasing, was observed between index and activity towards n-3 fatty acids (**BLUE**).

In proportion, the  $\Delta 6D$  activity towards n-3 fatty acids of slow-growing rabbits was higher respect that of the commercial Genotype (64.8 vs 10.9%, respectively).

**Table 1.** Estimated and enzyme activity of  $\Delta 6$ -desaturase (pmol in 30 min/mg protein) in rabbits fed different diets or belonging to different genotypes.

	Diets		Pooled SE <sup>1</sup>
	Control	Linolenic	
Estimated $\Delta 6$ -desaturase index (liver)	36.2 <sup>b</sup>	<b>28.9<sup>a</sup></b>	2.4
Real $\Delta 6$ -desaturase for C18:3n-3 activity	20.5 <sup>b</sup>	<b>17.2<sup>a</sup></b>	3.8
Real $\Delta 6$ -desaturase for C18:2n-6 activity	177.5	181.2	10.8
	Genotypes		Pooled SE <sup>1</sup>
	Commercial	Local	
Estimated $\Delta 6$ -desaturase index (liver)	31.8 <sup>a</sup>	<b>33.1<sup>b</sup></b>	1.5
Real $\Delta 6$ -desaturase for C18:3n-3 activity	19.1 <sup>a</sup>	<b>26.9<sup>b</sup></b>	2.4
Real $\Delta 6$ -desaturase for C18:2n-6 activity	175.2 <sup>b</sup>	41.5 <sup>a</sup>	15.0

Means in the same row and tissue with different letters differ significantly ( $P \leq 0.05$ ).

**Take home message:** the  $\Delta 6D$  index seems **usable** only towards the **activity of the n-3 series**; on the contrary, its trend and the total and n-6 real enzyme activities were not superimposable.