



## PROCEEDINGS OF THE 12<sup>th</sup> WORLD RABBIT CONGRESS

Nantes (France) - November 3-5, 2021

ISSN 2308-1910

### Session **PATHOLOGY & HYGIENE**

***Dakouri S.A. , Kimsé M., Koné M.W., Komoin O.C., Touré Alassane***

**EFFECTS OF *FICUS EXASPERATA*, *AZADIRACHTA INDICA* AND *MANGIFERA INDICA* LEAVES ON OOCYSTAL COCCIDIA EXCRETION AND ON RABBIT GROWTH**

**Full text of the communication**

**+**

**Slides of the oral presentation**

#### *How to cite this paper*

Dakouri S.A., Kimsé M., Koné M.W., Komoin O.C., Touré Alassane, 2021 Effects of *Ficus exasperata*, *Azadirachta indica* and *Mangifera indica* leaves on oocystal coccidia excretion and on rabbit growth. Proceedings 12th World Rabbit Congress - November 3-5 2021 - Nantes, France, Communication P-16, 4 pp. + presentation

## EFFECTS OF *FICUS EXASPERATA*, *AZADIRACHTA INDICA* AND *MANGIFERA INDICA* LEAVES ON OOCYSTAL COCCIDIA EXCRETION AND ON RABBIT GROWTH

Dakouri S. A<sup>1</sup>., Kimsé M<sup>1\*</sup>., Koné M. W.<sup>1</sup>, Komoin O. C.<sup>2</sup>, Touré Alassane<sup>2</sup>

1. Pôle Production Animales, UFR Sciences de la Nature, Université Nangui Abrogoua, 02 BP 801 Abidjan 02, Côte d'Ivoire

2. Laboratoire Central Vétérinaire de Bingerville (LCVB), Côte d'Ivoire.

\*Corresponding author: kimsemou.sn@univ-na.ci

### ABSTRACT

Synthetic anticoccidial curative action was decreased significantly. The aim of this study was to find an alternative cure to synthetic anticoccidials for the growing rabbits. We used 54 growing rabbits (6 groups and 9 rabbits per group). Three experimental groups (A, B and C) were used to test the effect of tropical leaves on coccidian. In addition, other 3 groups (T1, T2 and T3) were used to test the effect of sulfadimethoxine. Groups A, B and C diet was supplemented with *Ficus exasperata*, *Azadirachta indica* and *Mangifera indica* fresh leaves respectively. All animal of 3 groups were artificially infected with a coccidian inoculum at 25 days age. Group T1 was infected such as leaves groups and untreated; group T2 was not infected but it was treated with sulfadimethoxine and group T3 neither was infected nor received sulfadimethoxine. The mortality was low in group A (0 rabbits) compared to groups B, T3 (3 rabbits), C (5 rabbits) and T1 (9 rabbits) ( $P < 0.1$ ). It was similar to group T2 (2 rabbits). The secreted oocyst decreased in group A, C and T2 to 57%, 31% and 44% respectively from 35 to 65 days of age ( $P < 0.01$ ). However, oocyst rate was not reduced in group B, T1 and T3. It was increased (36%) in group T1 ( $P < 0.01$ ). *F. exasperata* leaves could be used as alternative to synthetic anticoccidials treatment in traditional rabbit breeding.

**Keywords:** Rabbit, Coccidian oocysts, tropical forage

### INTRODUCTION

Fattening rabbits are frequently confronted with digestive pathologies such as coccidiosis disease (Thoto, 2006). To fight against coccidiosis, the rabbit breeder generally uses sulfonamides or sulfadimethoxines in drinking water or rabbit food. However, the presence of sulfonamide residues in rabbit meat could affect the health of the consumer. Indeed, the body can develop resistance to antibiotics and toxicity (Kim and Lee, 2003). The emergence of antibiotic resistant bacteria is a major health problem. However, there are alternative solutions with the use of certain plants (Dossou, 2008). Several plants such as *F. exasperata* (Bafor *et al.*, 2009) and *M. indica* (Sereme *et al.*, 2008) are already used to treat digestive disorders in humans. The objective of this study is to reduce the effect of coccidia in rabbits using tropical plants.

### MATERIALS AND METHODS

The study was carried out in Côte d'Ivoire and the samples have been collected in Bingerville which is an endemic area (Kimsé *et al.*, 2016). Rabbit feces analyzes for the number of oocysts were carried out at the national laboratory for support to agricultural development (LANADA). Mortality was observed by counting animals that died of digestive disorders.

#### Animals and experimental devices

The feces of 54 rabbits (*Oryctolagus cuniculus*) weaned at 25 days of age was used. Rabbits were housed in individual cages. They were randomly divided into 6 groups (A, B, C, T1, T2 and T3). All animals received 120 g industrial food (FACI, Table 1). Group A, B and C were provided fresh leaves as additional to the industrial pellet. Group A, B and C received 150 g *Ficus exasperata*, *Azadirachta indica* and *Mangifera indica* fresh leaves respectively. Group

T1, T2 and T3 were fed with industrial pellet only. For coccidian effect, all animals (group A, B, C, T1 and T2) were infected artificially with coccidian except group T3. The group T1 was the positive control and T3 the negative control. To compare the effect of leaves to the synthetic treatment, group T2 received sulfadimethoxine 100-CR<sup>®</sup> (50 mg sulfadimethoxin per kg bodyweight per day for 30 days)

**Table 1.** Chemical composition the commercial feed

Nutrients	level (%)	Additifs
Crude protein	16.6	Vit. A 7500 U.I./Kg
Crude fat	3.9	Vit. D3 3000 U.I./Kg
Crude fiber	13.2	Vit. E 15 mg/Kg
Mineral matter	9.4	Vit. B1, 2, 3, 6, 12
Calcium	0.9	Lysine, Méthionine
Phosphate	0.9	Mg, Mn, Zn, Cu, Fe, Se
Sodium	0.3	

according to Mac Master method as described by Kimsé et al, (2016). The effect of treatment (leaves and sulfamid) was observed using a 3 g sample of droppings from each cage every 2 days.

### Statistical analyzes

Effects of treatment on the OPG number and their evolution during rabbit growth have been evaluated by an ANOVA 1 analysis at the threshold  $\alpha = 0.05$ . The comparison of means, in order to assess the impact of the treatment on animals, was carried out using the Tukey test. Analyzes were carried out using RStudio 3.1 software. The effect of treatment on mortality was evaluated in pairs using the chi-2 test at the threshold of  $\alpha = 0.05$  with excels program.

## RESULTS AND DISCUSSION

The highest mortality rate was observed on infected and untreated rabbits (group T1,  $P < 0.05$ ). All the rabbits in this group died (Table 2). Contrary, group A mortality (0%) was the lowest in this study ( $P < 0.05$ ). Nevertheless, group A was not differed significantly than group T2. However, groups T3 and C lost more animals than half their initial strength (5/9). This rate (56%) was not different to mortality rate in group B (34%) and T2 (23%). Group B and group T2 mortalities were not different.

**Table 2:** effect leaves supplementation and sulfadimethoxin on mortality

	Group A	Group B	Group C	Group T1	Group T2	Group T3
<i>Mortality</i>	0 (0)	3 (34)	5 (56)	9 (100)	2 (23)	5 (56)
Group A		0,06	0,01	0,01	0,13	0,01
Group B	0,06		0,34	0,01	0,60	0,34
Group C	0,01	0,34		0,02	0,14	1
Group T1	0,01	0,01	0,02		0,01	0,01
Group T2	0,13	0,60	0,14	0,01		0,14
Group T3	0,01	0,34	1	0,01	0,14	

() percentage of dead rabbit; Group A = infected rabbits whose diet was supplemented with *Ficus exasperata* leaves. Group B = infected rabbits whose diet was supplemented with *Azadirachta indica* leaves. Group C = infected rabbits whose diet was supplemented *Mangifera indica* leaves, Group T1 = infected rabbit and untreated, Group T2 = not infected rabbits and treated with sulfadimethoxin, Group T3 = not infected rabbits and untreated

The high mortality observed in this work was due to the high pathogenic of coccidia species encountered in this area. Thus, 11 species of coccidia were identified, including *E. magna*, *E. perforans*, *E. intestinalis* and *E. flavescens* which are very pathogenic (Ming-Hsien *et al.*, 2010; Kimsé *et al.*, 2016). This is specific to humid tropical areas. These areas are very favorable for the development of these digestive parasites (Akpo *et al.*, 2011). Without effective treatment, mortality increases when rabbits were infected with these highly virulent


species. In this study, mortality was low when the infected rabbit diet was supplemented with *F. exasperata* and *A. indica* fresh leaves. It was the same for infected rabbit treated with

sulfadimethoxin. These leaves could have a coccidiostatic action. The effect of the leaves of *F. exasperata* on coccidian was greater than *A. indica* and sulfadimethoxin. Which explains the sharp drop in the number of oocysts in *F. exasperata* (Table 3). Thus, the number of oocysts from rabbits treated with *F. exasperata* leaves reduced the infection. This forage could consistently maintain the number of oocysts as the sulfadimethoxin treatment. For the other groups, the peak of oocyst excretion was observed approximately 2 weeks after weaning (figure). This period would be the most favorable for preventive treatment because the rabbit's immune system is fragile.

**Table 3:** Effect of leaves on oocyst excretion during rabbit growth

	Ages (Days)				Standard deviation	P value
	35-42d	43-49d	50-57d	58-65d		
<i>Rabbit diet supplemented with fresh leaves</i>						
Lot A	19613 <sup>a</sup>	14663 <sup>b</sup>	10050 <sup>c</sup>	8350 <sup>c</sup>	3759	<0.01
Lot B	59500 <sup>c</sup>	103988 <sup>a</sup>	60500 <sup>b</sup>	42963 <sup>c</sup>	52588	<0.01
Lot C	44175 <sup>b</sup>	131500 <sup>a</sup>	32150 <sup>c</sup>	30450 <sup>c</sup>	52337	<0.01
<i>Rabbit diet not supplemented with fresh leaves</i>						
Lot T1	62338 <sup>c</sup>	186000 <sup>a</sup>	118013 <sup>b</sup>	97613 <sup>b</sup>	52031	<0.01
Lot T2	5050 <sup>a</sup>	4987 <sup>a</sup>	3265 <sup>b</sup>	2775 <sup>c</sup>	25667	<0.01
Lot T3	48375 <sup>c</sup>	100525 <sup>a</sup>	56788 <sup>b</sup>	50738 <sup>c</sup>	23282	<0.01

a, b, c: For the same line and for the same factor, the values followed by the same letter are not significantly different at the 5% threshold; Group A = infected rabbits whose diet was supplemented with *Ficus exasperata* leaves. Group B = infected rabbits whose diet was supplemented with *Azadirachta indica* leaves. Group C = infected rabbits whose diet was supplemented *Mangifera indica* leaves, Group T1 = infected rabbit and untreated, Group T2 = not infected rabbits and treated with sulfadimethoxin, Group T3 = not infected rabbits and untreated



**Figure:** Effect of treatment on oocysts excreted by rabbits from 35th to 65th day of age  
Group A = infected rabbits whose diet was supplemented with *Ficus exasperata* leaves. Group B = infected rabbits whose diet was supplemented with *Azadirachta indica* leaves. Group C = infected rabbits whose diet was supplemented *Mangifera indica* leaves, Group T1 = infected rabbit and untreated, Group T2 = not infected rabbits and treated with sulfadimethoxin, Group T3 = not infected rabbits and untreated

## CONCLUSION

The aim of this study was to propose an alternative to synthetic anticoccidials in breeding. Adding of *Ficus exasperata* and *Azadirachta indica* leaves to the diet reduced the mortality of growing rabbits unlike sulfadimethoxine. *F. exasperata* also reduced the number of oocysts excreted by more than half, better than sulfadimethoxine. The leaf of *F. exasperata* is a good alternative to synthetic anticoccidials for growing rabbits. However, additional studies are needed on a large population including growing rabbits and does before its widespread use.

## REFERENCES

- Akpo Y., Kpodekon M.T., Djago Y., Licois D., , Youssao I.A.K. 2011. Evaluation de l'innocuité des souches précoces de *Eimeria magna* et de *Eimeria media* issues du Bénin en vue de leur utilisation comme souches vaccinales. *Int. J.Biol. Chem.Sci.* 5 : 1682-1687.
- Bafor E.E., Igbinuwen O., 2009. Acute toxicity studies of the leaf extract of *Ficus exasperata* on haematological parameters, body weight and body temperature. *J. Ethnopharmacol*, 123, 302-307
- Dossou A.D., 2008. Effet du tourteau de Neem (*Azadirachta indica*. Juss) sur les coccidioses aviaires. *Thèse: Méd. Vét ; Dakar (Sénégal)*, 27p.
- Kimsé M., Dakouri S. A., Koné M. W., Komoin O. C., Coulibaly M., Yapi Y.M., Fantodji A.T., Otchoumou A. 2016. Rabbit's coccidian species in a tropical endemic area. *Proceedings - 11th World Rabbit Congress - June-15-18, 2016 – Qingdao, China*, 541-544
- MingHsien L.; Huang H.; HongKean O., 2010. Prevalence, infectivity and oocyst sporulation time of rabbit-coccidia in Taiwan. *Tropical biomedicine* 27(3):424-9
- Thoto M.C., 2006. Utilisation de la Robénidine (Cycostat ND 66G) en qualité d'additif anticoccidien dans l'aliment : effet sur la croissance et le degré d'infestation des lapins à l'engraissement. *Thèse méd. vét. Dakar (Sénégal)*, 5: 65pp



## EFFECTS OF FICUS EXASPERATA, AZADIRACHTA INDICA AND MANGIFERA INDICA LEAFS ON OOCYSTAL COCCIDIA EXCRETION AND ON RABBIT GROWTH



**Dakouri Serge Alain**

1. Context and issues
2. Objectives
3. Material and Methods
4. Results
5. Conclusion

# 1-Context and issues




Need  
+60%



Local  
Production

Solution



Coccidiosis



RESISTANCE

Alternative







**Reduce the effect of coccidia  
in rabbits using tropical  
plants**

# 3-Material and Methods

## Group design

- Group A = infected rabbits, diet supplemented with *Ficus exasperata* leaves
- Group B = infected rabbits, diet supplemented with *Azadirachta indica* leaves
- Group C = infected rabbits, diet supplemented with *Mangifera indica* leaves
- Group T1 = infected rabbit and untreated
- Group T2 = not infected rabbits and treated with sulfadimethoxine
- Group T3 = not infected rabbits and untreated



*Azadirachta indica*




*Mangifera indica*



*Ficus exasperata*

## A- Mc. Master method ( diagnosis and OPG)



## B- Culture + Floating test ( Identification)




# 4-Results

**Table 1: effect leaves supplementation and sulfadimethoxine on mortality**

	Group A	Group B	Group C	Group T1	Group T2	Group T3
<b>Mortality</b>	0 (0)	3 (34)	5 (56)	9 (100)	2 (23)	5 (56)
<b>Group A</b>		0,06	0,01	0,01	0,13	0,01
<b>Group B</b>	0,06		0,34	0,01	0,60	0,34
<b>Group C</b>	0,01	0,34		0,02	0,14	1
<b>Group T1</b>	0,01	0,01	0,02		0,01	0,01
<b>Group T2</b>	0,13	0,60	0,14	0,01		0,14
<b>Group T3</b>	0,01	0,34	1	0,01	0,14	

❖ In this study, mortality was low when the infected rabbit diet was supplemented with *F. exasperata* and *A. indica* fresh leaves. It was the same for infected rabbit treated with sulfadimethoxin (Tab 1).



**Figure 1: Effect of treatment on OPG from 35th to 65th day of age**

❖ The OPG from rabbits treated with *F. exasperata* leaves reduced the infection (Figure 1).

**Table 2: Effect of leaves on oocyst excretion during rabbit growth**

	Ages (Days)				Standard deviation	P value
	35-42d	43-49d	50-57d	58-65d		
Rabbit diet supplemented with fresh leaves						
Lot A	19613 <sup>a</sup>	14663 <sup>b</sup>	10050 <sup>c</sup>	8350 <sup>c</sup>	3759	<0.01
Lot B	59500 <sup>c</sup>	103988 <sup>a</sup>	60500 <sup>b</sup>	42963 <sup>c</sup>	52588	<0.01
Lot C	44175 <sup>b</sup>	131500 <sup>a</sup>	32150 <sup>c</sup>	30450 <sup>c</sup>	52337	<0.01
Rabbit diet not supplemented with fresh leaves						
Lot T1	62338 <sup>c</sup>	186000 <sup>a</sup>	118013 <sup>b</sup>	97613 <sup>b</sup>	52031	<0.01
Lot T2	5050 <sup>a</sup>	4987 <sup>a</sup>	3265 <sup>b</sup>	2775 <sup>c</sup>	25667	<0.01
Lot T3	48375 <sup>c</sup>	100525 <sup>a</sup>	56788 <sup>b</sup>	50738 <sup>c</sup>	23282	<0.01

❖ The effect of the leaves of *F. exasperata* on coccidian was greater than *A. indica* and sulfadimethoxin. Which explains the sharp drop in the number of oocysts in *F. exasperata* (Tab 2)

The aim of this study was to propose an alternative to synthetic anticoccidials in breeding. Adding *Ficus exasperata* and *Azadirachta indica* leaves to the diet reduced the mortality of growing rabbits unlike sulfadimethoxine. *F. exasperata* also reduced the number of oocysts excreted by more than half, better than sulfadimethoxine. The leaf of *F. exasperata* is a good alternative to synthetic anticoccidials for growing rabbits. However, additional studies are needed on a large population including growing rabbits and does before its widespread use.



***The authors thank Ivorian Rabbit Science Association (IRSA), C2D, AFD, FCIAD, FIRCA and GIDENNE T for their assistance in this study.***

